

Memorandum

April 30, 2021

To: Kristen Keene, Maryland Department of Transportation Maryland Port Administration
Cassandra Carr, Maryland Environmental Service

From: Karin Olsen, PG, Anchor QEA, LLC

Re: Elk River Sampling – River Beach Samples, Fall 2020

Program Overview

On behalf of the Maryland Department of Transportation Maryland Port Administration and the Maryland Environmental Service, sampling was conducted at two River Beach locations in the nearshore Elk River to assess the environmental conditions in the vicinity of the Pearce Creek Dredged Material Containment Facility (DMCF) Exterior Monitoring Area (Figure 1). The River Beach sampling efforts were initiated in 2016 based on environmental concerns expressed by citizen members of the Pearce Creek Implementation Committee. The purpose of this Memorandum is to summarize the results of the fall 2020 sediment quality characterization, water quality characterization, benthic community sampling, and benthic bioassay sampling for each of the two locations. Sample locations are shown on Figure 2 and coordinates are provided in Table 1.

Technical Approach

The data collection and analytical approach for the River Beach locations was consistent with the Pearce Creek DMCF Exterior Monitoring Program (Anchor QEA 2016a, 2016b, 2017a, 2017b, 2018, 2019a, 2019b, and 2020). The River Beach samples function as a discrete sample set and are evaluated independently from the samples collected in conjunction with the Pearce Creek DMCF Exterior Monitoring Program. Data collected during previous sampling events in spring and fall 2016, spring 2017, spring and fall 2018, and spring and fall 2019 are presented on the results tables (Tables 2 through 5) for comparison to data collected during the fall 2020 sampling event. The 2020 sampling event was conducted on October 5 and 6, 2020.

Sediment Quality Characterization

Undisturbed sediments were collected from the sediment-water interface to a depth of 6 inches using a Ponar grab sampler. Samples were submitted for metals, grain size, moisture content, specific gravity, total organic carbon (TOC), nitrate + nitrite, total Kjeldahl nitrogen (TKN), ammonia, total phosphorus, and sulfide. Chemical concentrations in bulk sediment samples were compared to sediment quality guidelines for freshwater samples (MacDonald et al. 2000).

Water Quality Monitoring

Surface water samples were collected from the mid-depth of the water column. Samples were submitted for dissolved metals, total suspended solids, phosphorus, hardness, ammonia, nitrate, and TKN analysis. Physical parameters, including temperature, dissolved oxygen, pH, and salinity, were also recorded at each sampling location. Chemical concentrations in the surface water samples were compared to the U.S. Environmental Protection Agency (USEPA) *National Recommended Water Quality Criteria* (2018) and the State of Maryland Code of Regulations (COMAR 26.08.02.03-2) freshwater acute water quality criteria for aquatic life.

Benthic Community Sampling

Benthic community (bottom-dwelling organisms) samples were collected to determine community composition, abundance (number of benthic organisms), and diversity (number of different types of species). The results were used to calculate benthic community metrics, including the number of total abundance, number of taxa, species richness, evenness, Shannon-Wiener Species Diversity Index, Simpson's Dominance Index, percent abundance of pollution indicative species, percent abundance of deep deposit feeders, and tolerance score.

Benthic Bioassays

Sediment was submitted for benthic bioassay testing. Benthic bioassays were used to evaluate if the sediments were acutely toxic to organisms living in the sediments. Bioassays were 10-day whole sediment tests using the freshwater amphipod *Hyaella azteca*. Testing was conducted according to the USEPA's *Methods for Measuring the Toxicity and Bioaccumulation of Sediment Associated Contaminants with Freshwater Invertebrates* (USEPA 2000). *Hyaella azteca* survival data for the whole sediment bioassays were statistically compared to the survival data in control sediment. A control sediment is a non-impacted sediment sample that is used to evaluate the results of a test.

Field Investigation


The methods and procedures for the collection of field samples, sampling schedule, rationale for the sampling design, and design assumptions for locating and selecting environmental samples were carried out in accordance with the Sampling and Analysis Plan (Anchor QEA 2015) and the methods used for the Pearce Creek DMCF Exterior Monitoring Program (Anchor QEA 2016a, 2016b, 2017a, 2017b, 2018, 2019a, 2019b, 2020). Sampling procedures were consistent with USEPA protocols or other approved sample collection standards. A complete list of analytes, target detection limits, and analytical methodologies is provided in the Sampling and Analysis Plan (Anchor QEA 2015).

Two River Beach (RB) sampling locations were included in this investigation. One location was near the dredged material inflow location for the Pearce Creek DMCF (location RB-01), and one location (location RB-02) was located approximately 1/3 mile downstream of RB-01. Sampling locations were

determined in the field using a Trimble ProXRS Differential Global Positioning System with an accuracy of 1 to 3 meters. Northing and easting coordinates for the sampling locations are provided in Table 1 and shown in Figure 2.

Sediment Quality Characterization

Concentrations of detected analytes in sediment samples were compared to consensus-based sediment quality guidelines for freshwater sediment, where available (MacDonald et al. 2000). Threshold effect concentrations (TECs) and probable effect concentrations (PECs) are derived based on empirical data from laboratory and field studies (MacDonald et al. 2000). The TEC values represent concentrations below which adverse biological effects are unlikely, and PEC values represent concentrations above which adverse biological effects are probable (MacDonald et al. 2000). Concentrations that are between the TEC and PEC represent the concentrations at which adverse biological effects might occur, as shown below:



Results of the sediment quality characterization are summarized in Table 2. In fall 2020, sample RB-01 was composed of 96% sand and 4% silts and clays. Sample RB-02 was composed of 18.7% gravel, 79.8% sand, and 1.5% silts and clays. TOC and nutrient concentrations were low at both locations. Nitrate + nitrite was detected at a concentration of 0.68 milligrams per kilogram (mg/kg) at RB-01 and 1.1 mg/kg at RB-02. TKN concentrations at RB-01 and RB-02 were 200 and 160 mg/kg, respectively. Ammonia was not detected in either sample. Total phosphorus was 42 mg/kg at RB-01 and 18 mg/kg at RB-02. Sulfide was detected at a concentration of 16 mg/kg at RB-01 but was not detected at RB-02. Nutrient concentrations in the fall 2020 sampling event were all within the range of the seven previous sampling events (spring 2016, fall 2016, spring 2017, spring 2018, fall 2018, spring 2019, and fall 2019).

Of the thirteen tested metals, nine were detected in at least one sample. Mercury, selenium, silver, and thallium were not detected at either location. Metal concentrations at both locations were low

and well below the TECs. Concentrations in both samples generally fell within the range of, or less than, concentrations reported in the previous sampling events (Table 2).

Water Quality Characterization

Analytes detected in the surface water were compared to the USEPA and State of Maryland freshwater acute and chronic water quality criteria. Criteria were derived from the USEPA *National Recommended Water Quality Criteria* (USEPA 2018) and the Code of Maryland Regulations (COMAR 26.08.02.03-2). For dissolved metals, the State of Maryland freshwater water quality criteria for the protection of aquatic life are the same as the USEPA criteria (Table 3) and are directly comparable to the results.

The State of Maryland allows, but does not require, that freshwater criteria be adjusted based on water hardness. The freshwater water quality criteria for the protection of aquatic life for cadmium, chromium, copper, lead, nickel, and zinc were calculated using the minimum hardness value (880 milligrams per liter [mg/L]), which was applied to both samples as a conservative evaluation of water quality.

Results of the water quality characterization are summarized in Table 3. Hardness values were similar between both samples (880 mg/L at RB-01 and 930 mg/L at RB-02). Nutrients were reported at similar concentrations between both surface water samples. Ammonia was not detected at either location. Total phosphorus was detected at a concentration of 0.046 mg/L at RB-01 and was not detected at RB-02. TKN was detected at concentrations of 2.2 mg/L and 2.8 mg/L at RB-01 and RB-02, respectively. The nitrate + nitrite concentration was 0.61 mg/L at RB-01 and 0.57 mg/L at RB-02. The total suspended solids concentration was 11 mg/L at RB-01 and 14 mg/L at RB-02. Nutrient concentrations in the fall 2020 sampling event generally fall within the range of concentrations from the previous seven sampling events (spring and fall 2016, spring 2017, spring and fall 2018, and spring and fall 2019) at RB-01 and RB-02.

Of the sixteen tested metals, seven were detected in one or both surface water samples (aluminum, antimony, arsenic, copper, iron, manganese, and nickel). None of the metals were detected at concentrations that exceeded acute or chronic freshwater criteria. Metal concentrations in both samples generally fell within the range of, or were less than, concentrations reported in the previous six sampling events (Table 3).

Benthic Community

Benthic (or bottom-dwelling) organisms are important indicators of stress in aquatic systems because they can integrate the effects of environmental conditions during long periods of time. Benthic organisms are also important food for many fish, providing an important link to higher trophic levels. Most benthic organisms tend to thrive only in some habitats (for example, sandy

versus muddy sediments), and groups of benthic organisms collected at sampling locations generally comprise species that are adapted to a specific habitat. Sampling locations are considered “normal” or “healthy” when the benthic organisms collected from that location are primarily the species that are specifically adapted to live in that particular habitat.

Results of the benthic community sampling are summarized in Table 4. The salinity measured at RB-01 and RB-02 was 3.4 parts per thousand (ppt) and 3.7 ppt, respectively (Table 1); therefore, both locations were classified as oligohaline (bottom salinity ranging from 0.5 to 5 ppt).

Total benthic abundance (total number of organisms per square meter [m^2]) was 1,367 organisms/ m^2 at RB-01 and 3,490 organisms/ m^2 at RB-02 (Table 4). Twenty-two benthic taxa were collected from the River Beach locations. Thirteen taxa were collected at RB-01: Diptera (five taxa), Polychaeta (four taxa), Oligochaeta (two taxa), and Isopoda (two taxa). Twenty taxa were collected at RB-02: Diptera (ten taxa), Polychaeta (five taxa), Oligochaeta (two taxa), Bivalvia (two taxa), and Isopoda (one taxon). Tubificidae was the dominant taxa at RB-01 and benthic organism abundance at RB-02 was dominated by Diptera (specifically Polypedilum).

Species richness is a comparison of how many taxa are in a sample compared to how many individuals are in a sample. Lower values indicate that the total benthic abundance at a location is dominated by a few taxa and does not represent a diverse benthic community. The species richness at RB-01 was 2.3 and the species richness at RB-02 was 2.7. Species richness values were comparable with values observed in previous years (Table 4).

Evenness is a measure of how evenly the individuals collected at a location are distributed among the taxa collected at that location, with a value of 1 indicating that the individuals are distributed as evenly as possible. Evenness values at RB-01 and RB-02 were 0.80 and 0.88, respectively. The evenness values at RB-01 and RB-02 were comparable to, or slightly greater than, those observed in all seven previous monitoring events (Table 4).

The Shannon-Wiener Species Diversity Index takes into account species richness and species evenness, with higher values indicating a more diverse benthic community. Location RB-01 and RB-02 had a Shannon-Wiener Species Diversity Indices of 2.8 and 3.4, respectively. The Shannon-Wiener Species Diversity Index measured at RB-01 was within the range of values observed in the previous monitoring events, however the Shannon-Wiener Species Diversity Index measured at RB-02 was greater than all values previously measured (Table 4).

Simpson’s Dominance Index measures the diversity of a sample, with a lower value indicating a more diverse community. Simpson’s Dominance Index was 0.2 at RB-01 and 0.12 at RB-02 (Table 4). The results for RB-01 were within the range of those observed during previous monitoring events;

however, the results for RB-02 were less than the values measured during all previous monitoring events.

Results for the benthic community evaluation for fall 2020 were generally consistent with the results for the seven previous sampling events (spring 2016, fall 2016, spring 2017, spring 2018, fall 2018, spring 2019, and fall 2019; Table 4). The benthic metrics were generally within the range of, or showed improvement upon, those observed in the previous seven sampling events (Table 4). This indicates that while the species composition of the benthic community changes seasonally in response to temperature, salinity, and dissolved oxygen fluctuations, the overall health of the benthic community is stable and possibly improving.

Benthic Bioassays

Benthic bioassays with whole sediment are designed to determine whether the sediment from each sampling location is likely to produce unacceptable adverse effects on benthic organisms by exposing the organisms to the whole sediment for 10 days. A freshwater amphipod (*Hyaletta azteca*) was used in the whole-sediment bioassay.

Hyaletta azteca is adapted to live in silty environments, so the toxicity tests are only applicable for fine-grained sediments comprising mostly silts and clays. However, for the fall 2020 sampling event, both locations comprised primarily coarse-grained material—RB-01 was 96% sand and RB-02 was 98.5% sands and gravel. Even though the substrate at both locations was coarse-grained, bioassay was conducted on both River Beach locations to evaluate site conditions for benthic organisms.

Results of the benthic bioassays were compared to the results in the control (Table 5). A control sediment is a non-impacted sediment sample that is used to evaluate the results of a test. The mean survival result of *Hyaletta azteca* exposed for 10 days to the River Beach sediment sample locations was 100% and 98% at RB-01 and RB-02, respectively. The survival result was not statistically different ($p=0.05$) from the mean survival in the control sediment (94%). Therefore, the sediment sample collected from location RB-01 and RB-02 was unlikely to cause adverse effects to benthic organisms.

Benthic bioassay results for the fall 2020 samples were comparable with the results for spring and fall 2016, spring 2017, spring and fall 2018, and spring and fall 2019, with samples from each event indicating that the sediment samples collected from locations RB-01 and RB-02 are unlikely to cause adverse effects to benthic organisms.

Summary

Sampling was conducted for two River Beach locations in the nearshore Elk River to evaluate existing conditions for sediment quality, surface water quality, benthic community, and benthic bioassays. Data collected during this investigation was compared to the previous sampling events (spring and

fall 2016, spring 2017, spring and fall 2018, and spring and fall 2019) and will be compared to any potential future data collection efforts to identify any trends or changes in sediment quality, surface water quality, benthic community, and benthic bioassays. The data collected over the course of this monitoring program will be analyzed and used to determine the need for additional monitoring events in the future.

References

Anchor QEA (Anchor QEA, LLC), 2015. *Sampling and Analysis Plan, Pearce Creek Dredged Material Containment Facility Exterior Monitoring Program*. Prepared for Maryland Environmental Service. September.

Anchor QEA, 2016a. *Pearce Creek Dredged Material Containment Facility Exterior Monitoring Program: Monitoring Report – Fall 2015*. Prepared for Maryland Port Administration and Maryland Environmental Service. March.

Anchor QEA, 2016b. *Pearce Creek Dredged Material Containment Facility Exterior Monitoring Program: Monitoring Report – Spring 2016*. Prepared for Maryland Port Administration and Maryland Environmental Service. October.

Anchor QEA, 2017a. *Pearce Creek Dredged Material Containment Facility Exterior Monitoring Program: Monitoring Report – Fall 2016*. Prepared for Maryland Port Administration and Maryland Environmental Service. April 2017.

Anchor QEA, 2017b. *Pearce Creek Dredged Material Containment Facility Exterior Monitoring Program: Monitoring Report – Spring 2017*. Prepared for Maryland Port Administration and Maryland Environmental Service. December 2017.

Anchor QEA, 2018. *Pearce Creek Dredged Material Containment Facility Exterior Monitoring Program: Monitoring Report – Spring 2018*. Prepared for Maryland Port Administration and Maryland Environmental Service.

Anchor QEA, 2019a. *Pearce Creek Dredged Material Containment Facility Exterior Monitoring Program: Monitoring Report – Fall 2018*. Prepared for Maryland Port Administration and Maryland Environmental Service.

Anchor QEA, 2019b. *Pearce Creek Dredged Material Containment Facility Exterior Monitoring Program: Monitoring Report – Spring 2019*. Prepared for Maryland Port Administration and Maryland Environmental Service.

Anchor QEA, 2020. *Pearce Creek Dredged Material Containment Facility Exterior Monitoring Program: Monitoring Report – Fall 2019*. Prepared for Maryland Port Administration and Maryland Environmental Service.

MacDonald, D.D., C.G. Ingersoll, and T.A. Berger, 2000. Development and Evaluation of Consensus-Based Sediment Quality Guidelines for Freshwater Ecosystems. *Archives of Environmental Contamination and Toxicology* 39:20-31.

USEPA (U.S. Environmental Protection Agency), 2000. *Methods for Measuring the Toxicity and Bioaccumulation of Sediment Associated Contaminants with Freshwater Invertebrates, Second Edition*. EPA 600/R-99/064. March 2000.

USEPA, 2018. National Recommended Water Quality Criteria. Accessed: August 28, 2018. Available at: <http://water.epa.gov/scitech/swguidance/standards/criteria/current/index.cfm>.

Tables

Table 1
Sample Collection and Water Quality Parameters

Location	Date	Time (EST)	Northing ^a	Easting ^a	Water Depth (feet)	Temperature (°C)	Salinity (ppt)	Dissolved Oxygen (mg/L)	Turbidity (NTU)	pH
RB-01	10/6/2020	1124	645821.28	1599391.2	5.5	19.0	3.4	8.6	7.0	7.8
RB-02	10/6/2020	1022	645053.15	1598004.6	5	18.7	3.7	9.9	4.8	7.9

Notes:

a: Coordinates are in Maryland State Plane, North American Datum of 1983.

EST: Eastern Standard Time

mg/L: milligram per liter

NTU: Nephelometric Turbidity Unit

ppt: part per thousand

Sample data recorded from middle depth location.

Table 2
Analytical Results for Sediment Samples

Analyte	Units	TEC	PEC	River Beach Location 1							
				Spring 2016	Fall 2016	Spring 2017	Spring 2018	Fall 2018	Spring 2019	Fall 2019	Fall 2020
Physical Characteristics											
Gravel	%	--	--	9.4	40.4	1.4	0.8	0	2.2	1.5	0.01 U
Sand	%	--	--	20.7	59	97	97.2	97.3	96.5	96.2	96
Silt	%	--	--	37	0.4	0.02	0.4	1.1	0.01 U	0.5	2.4
Clay	%	--	--	32.9	0.2	1.6	1.6	0.5	1.3	1.8	1.6
Specific Gravity	--	--	--	2.64	2.67	2.68	2.67	2.68	2.68	2.67	2.66
Nutrients											
Total Organic Carbon	%	--	--	2.9	0.17	0.62	0.33	0.19	0.14 U	0.14 U	0.42
Nitrate + Nitrite	mg/kg	--	--	4.2	1.3 U	1.3 J	1.5	1.6	1.2 J	1.7	0.68 J
Total Kjeldahl Nitrogen	mg/kg	--	--	2,200	140 J	390 U	200 J	150 J	160 J	130 J	200 J
Ammonia	mg/kg	--	--	150	10	20	8.9 J	7.8 U	9.5 J	11 J	14 U
Total Phosphorus	mg/kg	--	--	620	31	78	51	58	49	14	42
Sulfide	mg/kg	--	--	460	38 U	73 U	25 J	41 U	16 J	41 U	16 J
Metals											
Antimony	mg/kg	--	--	0.29	0.11 J	0.11 J	0.30	0.077 J	0.062 J	0.14 U	0.037 J
Arsenic	mg/kg	9.79	33	7.1	1.9	1.3	1.1	1.0	0.7	0.5	0.82
Beryllium	mg/kg	--	--	1.3	0.4	0.21	0.14	0.1	0.1	0.13	0.1
Cadmium	mg/kg	0.99	4.98	0.31	0.21	0.043 J	0.042 J	0.055 J	0.021 J	0.023 J	0.033 J
Chromium	mg/kg	43.4	111	29	7.4	8.6	5.7	6.3	6.0	5.9	5.8
Copper	mg/kg	31.6	149	21	1.8	2.3	1.8	3.3	1.6	1.3	1.7
Lead	mg/kg	35.8	128	32	1.5	5.1	5.1	3.7	3.5	2.2	2.4
Mercury	mg/kg	0.18	1.06	0.08	0.019 U	0.041 U	0.041 U	0.025 U	0.021 U	0.022 U	0.019 U
Nickel	mg/kg	22.7	48.6	33	3.1	4.1	4.1	2.7	2.6	2.2	3.1
Selenium	mg/kg	--	--	1.6	0.5	0.25 J	0.25 J	0.087 J	0.25 J	0.34 U	0.35 U
Silver	mg/kg	--	--	0.25	0.008 J	0.12 U	0.12 U	0.038 J	0.07 U	0.068 U	0.07 U
Thallium	mg/kg	--	--	0.15	0.0049 J	0.012 J	0.012 J	0.018 J	0.07 U	0.068 U	0.07 U
Zinc	mg/kg	121	459	120	13	19	9.7	11	10	6.8	12

Notes:

Bold indicates detected constituents.

 : constituents that exceed probable effect concentration

--: no value

J: estimated value; result is less than the reporting limit but greater than the method detection limit

mg/kg: milligram per kilogram

PEC: probable effects concentration

TEC: threshold effects concentration

U: compound not detected

Table 2
Analytical Results for Sediment Samples

Analyte	Units	TEC	PEC	River Beach Location 2							
				Spring 2016	Fall 2016	Spring 2017	Spring 2018	Fall 2018	Spring 2019	Fall 2019	Fall 2020
Physical Characteristics											
Gravel	%	--	--	7.8	17.0	9.6	15.1	17.4	5.4	7.9	18.7
Sand	%	--	--	91	81.5	87.1	84.1	76.3	93.5	88.8	79.8
Silt	%	--	--	0.4	0.9	1.7	0	5.1	0.4	2	0.03
Clay	%	--	--	0.8	0.6	1.6	0.8	1.3	0.8	1.3	1.5
Specific Gravity	--	--	--	2.69	2.66	2.67	2.67	2.65	2.68	2.68	2.67
Nutrients											
Total Organic Carbon	%	--	--	0.15	0.15	0.13 U	0.23	0.15	0.13 U	0.13 U	0.23
Nitrate + Nitrite	mg/kg	--	--	1.6	0.58 J	1.2 U	2	1.1 J	1.4	1.3	1.1 J
Total Kjeldahl Nitrogen	mg/kg	--	--	210	96 J	200 U	540	300	200	130 J	160 J
Ammonia	mg/kg	--	--	12 U	8.2	10.0	8.2 U	6.7 U	13 U	14 U	12 U
Total Phosphorus	mg/kg	--	--	42	31	30	33	17	24	18	18
Sulfide	mg/kg	--	--	9.8 J	9.1 J	38 U	22 J	36 U	39 U	40 U	38 U
Metals											
Antimony	mg/kg	--	--	0.077 J	0.05 J	0.029 J	0.061 J	0.053 J	0.13 U	0.077 J	0.032 J
Arsenic	mg/kg	9.79	33	0.82	0.50	0.47	0.45	0.57	0.65	0.64	0.49
Beryllium	mg/kg	--	--	0.08	0.059 J	0.054 J	0.066 J	0.18	0.036 J	0.15	0.058 J
Cadmium	mg/kg	0.99	4.98	0.013 J	0.21	0.017 J	0.014 J	0.029 J	0.012 J	0.016 J	0.019 J
Chromium	mg/kg	43.4	111	4.3	4.7	3.5	3.8	18	3.5	8.6	3.9
Copper	mg/kg	31.6	149	1.6	1.1	0.93	1.2	5.3	0.78	1.1	0.79
Lead	mg/kg	35.8	128	2	1.6	1.6	1.7	5.3	1.5	3.1	1.6
Mercury	mg/kg	0.18	1.06	0.0042 J	0.02 U	0.02 U	0.022 U	0.02 U	0.02 U	0.022 U	0.018 U
Nickel	mg/kg	22.7	48.6	1.4	1.1	1.2	1.4	2.5	1.4	1.4	1.3
Selenium	mg/kg	--	--	0.091 J	0.19 J	0.12 J	0.07 J	0.082 J	0.15 J	0.33 U	0.31 U
Silver	mg/kg	--	--	0.0053 J	0.008 J	0.063 U	0.071 U	0.061 U	0.064 U	0.066 U	0.062 U
Thallium	mg/kg	--	--	0.0063 J	0.0036 J	0.0036 J	0.071 U	0.0083 J	0.064 U	0.066 U	0.062 U
Zinc	mg/kg	121	459	5.1	5.2	5.1	5.1	8.3	4.9	5.9	4.7

Notes:

Bold indicates detected constituents.

--: no value

J: estimated value; result is less than the reporting limit but greater than the method detection limit

mg/kg: milligram per kilogram

PEC: probable effects concentration

TEC: threshold effects concentration

U: compound not detected

Table 3
Analytical Results for Surface Water Samples

Analyte	Unit	Acute Water Quality Criteria	Chronic Water Quality Criteria	River Beach Location 1							
				Spring 2016	Fall 2016	Spring 2017	Spring 2018	Fall 2018	Spring 2019	Fall 2019	Fall 2020
Hardness	mg/L	--	--	86	880	72	86	70	60	64	880
Total Phosphorus	mg/L	--	--	0.049 J	0.14	0.1 U	0.11	0.051 J	0.1 U	0.1 U	0.046 J
Total Suspended Solids	mg/L	--	--	11	40	8.9	39	10	4	6.4	11
Ammonia	mg/L	--	--	0.2	0.21	0.18	0.38	0.12	0.048 J	0.11	0.1 U
Total Kjeldahl Nitrogen	mg/L	--	--	5 U	2.2 J	11	1.7 J	5 U	5 U	5 U	2.2 J
Nitrate+Nitrite	mg/L	--	--	0.85	0.41	0.66	0.69	1.2	1.4	0.9	0.61
Metals											
Aluminum	µg/L	750	87	19 J	33	30 U	190	67	30 U	19 J	30 U
Antimony	µg/L	--	--	0.27 J	0.61 J	1.5 J	2 U	2 U	0.39 J	0.41 J	0.39 J
Arsenic	µg/L	340	150	0.83 J	0.77 J	0.34 J	1.4	0.65 J	0.44 J	0.96 J	1
Beryllium	µg/L	--	--	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Cadmium ^a	µg/L	13.6	3.7	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Chromium ^a	µg/L	3,382	440	1.3 J	0.39 J	2 U	2.2	1.1 J	2 U	2 U	2 U
Copper ^a	µg/L	104	57	1.2 J	1.9 J	2 U	2	1.3 J	0.96 J	0.97 J	0.94 J
Iron	µg/L	--	1,000	31 J	88	50 U	460	120	50 U	25 J	50 U
Lead ^a	µg/L	617	24	1 U	0.25 J	1 U	0.38 J	0.14 J	1 U	1 U	1 U
Manganese	µg/L	--	--	3.9 J	810	5 U	260	15	2 J	40	3.3 J
Mercury	µg/L	1.40	0.77	0.2 U	0.2 U	0.2 U	0.2 U	0.2 U	0.2 U	0.2 U	0.2 U
Nickel ^a	µg/L	2,948	327	1.2	4.6	1	3.5	1.3	1	1.8	1.8
Selenium	µg/L	20	5	5 U	0.57 J	5 U	5 U	5 U	5 U	5 U	5 U
Silver ^a	µg/L	135	--	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Thallium	µg/L	--	--	1 U	1 U	0.054 J	1 U	1 U	1 U	1 U	1 U
Zinc ^a	µg/L	740	746	4.2 J	4.2 J	5 U	3.9 J	5.1	5 U	14	5 U

Notes:

Bold indicates detected constituents.

: constituents that exceed chronic criteria

a. Acute and chronic water quality criteria are adjusted for a hardness of 880 mg/L.

µg/L: microgram per liter

J: estimated value; result is less than the reporting limit but greater than the method detection limit

mg/L: milligram per liter

U: compound not detected

Table 3
Analytical Results for Surface Water Samples

Analyte	Unit	Acute Water Quality Criteria	Chronic Water Quality Criteria	River Beach Location 2							
				Spring 2016	Fall 2016	Spring 2017	Spring 2018	Fall 2018	Spring 2019	Fall 2019	Fall 2020
Hardness	mg/L	--	--	86	940	70	86	72	62	660	930
Total Phosphorus	mg/L	--	--	0.1 U	0.1	0.037 J	0.1 U	0.1 U	0.1 U	0.1 U	0.1 U
Total Suspended Solids	mg/L	--	--	8.4	22	7.1	29	6.1	5.4	7	14
Ammonia	mg/L	--	--	0.15	0.16	0.16	0.21	0.051 J	0.1 U	0.1 U	0.1 U
Total Kjeldahl Nitrogen	mg/L	--	--	5 U	2.2 J	3.4 J	5 U	5 U	5 U	5 U	2.8 J
Nitrate+Nitrite	mg/L	--	--	0.83	0.25	0.65	0.95	1.2	1.4	0.93	0.57
Metals											
Aluminum	µg/L	750	87	16	48	16 J	22 J	14 J	30 U	30 U	25 J
Antimony	µg/L	--	--	0.26 J	0.93 J	0.98 J	2 U	2 U	0.4 J	0.43 J	0.39 J
Arsenic	µg/L	340	150	0.77 J	1.3	0.41 J	1.2	0.69 J	0.47 J	0.99 J	1
Beryllium	µg/L	--	--	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Cadmium ^a	µg/L	13.6	3.7	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Chromium ^a	µg/L	3,382	440	1.2 J	0.55 J	2 U	1.9 J	1.1 J	2 U	2 U	2 U
Copper ^a	µg/L	104	57	1.3 J	2.4	2 U	1.7 J	1.4 J	1.1 J	0.97 J	0.95 J
Iron	µg/L	--	1,000	28 J	51	23 J	37 J	26 J	50 U	50 U	44 J
Lead ^a	µg/L	617	24	1 U	0.35 J	1 U	1 U	0.15 J	1 U	1 U	1 U
Manganese	µg/L	--	--	4 J	43	3.2 J	5.4	8.9	19	34	14
Mercury	µg/L	1.40	0.77	0.2 U	0.2 U	0.2 U	0.2 U	0.2 U	0.2 U	0.2 U	0.2 U
Nickel ^a	µg/L	2,948	327	1.2	2.6	0.69 J	1.6	1.3	1.2	1.6	1.6
Selenium	µg/L	20	5	5 U	0.96 J	5 U	5 U	5 U	5 U	5 U	5 U
Silver ^a	µg/L	135	--	1 U	0.3 J	1 U	1 U	1 U	1 U	1 U	1 U
Thallium	µg/L	--	--	1 U	1 U	1 U	1 U	1 U	1 U	1 U	1 U
Zinc ^a	µg/L	740	746	3.4 J	3.5 J	5 U	5 U	5.1	5 U	4.4 J	5 U

Notes

Bold indicates detected constituents.

a. Acute and chronic water quality criteria are adjusted for a hardness of 880 mg/L.

µg/L: microgram per liter

J: estimated value; result is less than the reporting limit but greater than the method detection limit

mg/L: milligram per liter

U: compound not detected

Table 4
Benthic Community Metrics

Metric	River Beach Location 1							
	Spring 2016	Fall 2016	Spring 2017	Spring 2018	Fall 2018	Spring 2019	Fall 2019	Fall 2020
Total Abundance/m ²	1,907	1,773	2,250	3,509	2,727	2,892	1,697	1,367
Infaunal Taxa	14	15	12	16	11	15	21	13
Species Richness (Ludwig-Reynolds)	2.6	3.1	2.3	2.9	2.0	2.8	4.0	2.3
Evenness	0.74	0.67	0.69	0.78	0.48	0.80	0.78	0.80
Shannon-Wiener H' (log base 2)	2.7	2.6	2.5	3.1	1.7	3.1	3.3	2.8
Simpson's Dominance Index	0.21	0.25	0.24	0.15	0.41	0.15	0.14	0.20
Percent Abundance Pollution Indicative Species	38	43	21	18	18	27	75	75
Percent Abundance Deep Deposit Feeders	38	0	33	45	19	46	35	47
Tolerance Score	5.05	1.30	5.6	5.8	5.6	6.7	6.3	5.7

Metric	River Beach Location 2							
	Spring 2016	Fall 2016	Spring 2017	Spring 2018	Fall 2018	Spring 2019	Fall 2019	Fall 2020
Total Abundance/m ²	2,333	3,502	2,981	7,024	7,462	11,066	2,117	3,490
Infaunal Taxa	15	12	11	12	16	18	14	20
Species Richness (Ludwig-Reynolds)	2.5	2.1	2.0	1.9	2.5	2.7	2.3	2.7
Evenness	0.73	0.68	0.76	0.77	0.42	0.49	0.68	0.88
Shannon-Wiener H' (log base 2)	2.7	2.4	2.6	2.8	1.7	2.1	2.4	3.4
Simpson's Dominance Index	0.21	0.24	0.20	0.19	0.42	0.18	0.30	0.12
Percent Abundance Pollution Indicative Species	32	66	14	3	26.9	37.7	88.0	51.4
Percent Abundance Deep Deposit Feeders	62	0	24	57	28.5	48.1	27.9	17.5
Tolerance Score	8.04	4.52	4.8	7.0	6.75	5.7	5.4	4.4

Note:

m²: square meter

Table 5
Summary of Test Acceptability Endpoints for Whole Sediment Acute Bioassay for *Hyalella azteca*

Endpoint/ Measurement	Protocol Criteria	Units	Spring 2016	Fall 2016	Spring 2017	Spring 2018	Fall 2018	Spring 2019	Fall 2019	Fall 2020
Survival	Mean Laboratory Control	Mean Survival %	94%	94%	94%	91%	91%	100%	94%	94% ^a
	≥ 80%	Protocol Met	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Growth	Measure Positive Growth	Start Dry Weight (mg)	0.024	0.017	0.018	0.008	0.0343	0.0258	0.0234	0.0365
	End vs. Start of Assay	End Dry Weight (mg)	0.143	0.124	0.147	0.659	0.102	0.134	0.0969	0.104
	Protocol Met		Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Temperature	Mean: 23 °C ± 1 °C	Daily/Hourly	22.8 / 22.8	21.3 / 21.6	23.3 / 23.4	22.0 / 21.9	22.9 / 20.2	23.4 / 23.4	22.6 / 22.7	22.8 / 22.9
	Minimum: 20 °C	Daily/Hourly	22.1 / 21.7	20.2 / 20.1	22.9 / 22.9	20.9 / 20.9	22.3 / 18.2	22.8 / 21.6	21.9 / 21.3	22.2 / 21.6
	Maximum: 26 °C	Daily/Hourly	23.4 / 23.4	22.4 / 22.5	23.6 / 23.9	22.5 / 23.2	23.3 / 20.9	24.2 / 24.8	23.2 / 23.4	23.7 / 23.3
	Protocol Met		Yes / Yes	No / Yes	Yes / Yes	Yes / Yes	Yes / No ^b	Yes / Yes	Yes / Yes	Yes / Yes

Note:

a. Mean *Hyalella azteca* survival was 100% at RB-01 and 98% at RB-02.



b. The hourly temperature measurements recorded for the assay fell below the acceptable thresholds required for the mean and minimum temperatures. However, daily temperature measurements were all within the acceptable range. This deviation had no adverse impact on the outcome of the assay.

mg: milligram

Figures




LEGEND:

-  Exterior Monitoring Area
-  Pearce Creek DMCF

NOTE:

1. Base map courtesy of Esri and its data suppliers (2017).



Publish Date: 2021/01/21, 9:03 AM | User: jqinley
 Filepath: \\orcas\GIS\Jobs\Maryland_ES_0530\PearceCreek_Exterior_Monitoring_SAP\Maps\MonitoringReports\Fall2020\MES_PearceCrk_VicinityMap.mxd



Figure 1
Site Location Map
 Elk River Sampling: Fall 2020
 Pearce Creek DMCF Exterior Monitoring Program




LEGEND:

- Exterior Monitoring Location
- Pearce Creek Dredged Material Containment Facility

NOTE:

1. Aerial imagery: NAIP 2017.



Publish Date: 2021/01/21, 9:06 AM | User: jquinley

Filepath: \\orcas\GIS\Jobs\Maryland_ES_0530\PearceCreek_Exterior_Monitoring_SAP\Maps\MonitoringReports\Fall2020\MES_ElkRiver_Beach_Sampling_Fall2020.mxd



Figure 2
Pearce Creek Beach Sampling Locations

Elk River Sampling: Fall 2020
Pearce Creek DMCF Exterior Monitoring Program